

Pediatric Surgery: A New Surgical Specialty in Nepal and its Critical Issues

Sushil Rijal*,
Fatima Naumeri and
Muhammad Sharif

King Edward Medical University, Lahore,
Pakistan

*Corresponding author: Sushil Rijal

✉ Sushil.rijal@gmail.com

King Edward Medical University, Lahore,
Pakistan.

Tel: +923334449639

Citation: Rijal S, Naumeri F, Sharif M (2021) Pediatric Surgery: A New Surgical Specialty in Nepal and its Critical Issues. *Pediatr Emerg Care Med Open Access*. Vol.6 No.2:9

Abstract

Practice of pediatric surgery started in mid-14th century when both medicine and surgery, as a whole science were at its best primal stage. As there was little knowledge about resuscitation of children and their anesthesia, naïveté of visceral lesions and its operative technique, operative practice in pediatric population was limited to deal with some of congenital or acquired orthopedic diseases, removal of urinary bladder stones or some minor surgeries.

Keywords: Orthopedic diseases; Urinary bladder; Resuscitation

Received: September 03, 2020; **Accepted:** March 17, 2021; **Published:** March 24, 2021

Introduction

Although, the history of pediatric surgery as a sub specialty is not that long [1,2]. Reported annual infant mortality is approximately 4.5 million worldwide, and 99% of it is contributed by developing countries. Recorded deaths of children below five years of age are around 10 million annually [3]. According to Data published by National Population and Housing Census Nepal 2011 and Nepal Demographic and Health Survey (NDHS) 2011, Nepal is the 49th most populous country in world counting more than 26 million people, among them 34.6% is below 14 years of age which falls under pediatric age group. Besides, Infant mortality rate (IMR) is 46 per 1000 live births and Under 5 mortality rate (U5MR) is 39 per 1000 live births which is higher than some of south Asian countries. IMR and U5MR are gradually dilapidating in last few decades globally but developing countries with low income like Nepal, is still lagging and the rate is above the Sustainable Development Goal target at present [4-7].

Literature Review

Nepal has very short history of modern health services where only in 1956 health ministry was established and control and prevention of communicable diseases was its main concern. Such high IMR and U5MR in Nepal are largely related with birth asphyxia, prematurity, sepsis and surgery related problems. In low and middle income developing countries Pediatric surgery is still a challenge, particularly in the emergency setup. The common surgical issues of children frequently faced in emergency are mostly trauma related and congenital abnormalities. Because of multifaceted and expensive treatment, lack of proper trained pediatric surgeons, lack of pediatric surgery setup like

postoperative wards and NICU, the desired functional outcome is not easy to achieve in developing countries which is why the mortality rate in postoperative neonate is 6.4% in a developed country and is 62.2% in a developing country [8-10].

Surgical procedure is globally very large. More than 2 billion people are expected to be beyond surgical care access. In countries with low or middle income, very little statistics on surgical conditions for pediatric populations are available. However one study done in African setup reveals 85% of children needs surgical consultation and treatment before they reach 15years. Our local study shows roughly seven hundred thousand children need surgical consultation in Nepal annually [11-13].

Pediatric surgery as a subspecialty is recognized in 87.5% of countries globally with 6.3 years mean duration of training. Those countries were barred which demands fully qualified general surgeon before doing pediatric surgery. Moreover, tissue engineering is also rising along with pediatric surgery as numerous tissue deficit diseases in pediatric surgical cases; treatment is being achieved through tissue engineering [14,15].

Nepal, being low-income developing country encounters significant hurdle in delivering surgical care in pediatric population. Facility of postgraduate training for surgeons/health care providers has not been available till date in whole country of 26 million populations. Currently, most of the pediatric surgical cases are taken up by general surgeons who have never

trained in pediatric surgery program. Also, majority of them lack experience achieved by academic pediatric surgeon. On the contrary, advanced pathology, late presentation, minimal pediatric surgeons within the country, lack of proper trained pediatric surgeons outside the tertiary hospitals and inadequate governmental support are the challenges Nepal is facing now.

Discussion and Conclusion

With the help general and orthopedic surgeons we have managed so far though it is not their principal domain. Now is the time to off load burden of our fellow general and orthopedic surgeons by introducing pediatric surgery as subspecialty and take over. To improve pediatric health standard and to provide better surgical consultation to large pediatric population of Nepal, immediate establishment of postgraduate training opportunities within country and alliance with well-established pediatric surgical training centers/universities of developed countries is mandatory.

Conflict of Interest

The authors declared no conflict of interest.

References

- 1 Chang JH (1986) Timelines in the history of pediatric surgery. *J Pediatr Surg* 21: 1068-1072.
- 2 Prévot J (1986) The history of pediatric surgery in France. In *Historical Aspects Pediatr Surg*, pp: 9-16.
- 3 Naumeri F, Rijal S, Awan NU, Farooq MS (2018) Emergency Neonatal Surgery: Outcome in a University Hospital of Pakistan. *Internat J Adv Biotechnol Res* 9: 123-128.
- 4 CBS News (2012) National population and housing census 2011. National Report.
- 5 Kathmandu N (2014) Central Bureau of Statistics. *Populat* 33: 34-40.
- 6 Ghimire PR, Agho KE, Ezeh OK, Renzaho A, Dibley M, et al. (2019) Under-five mortality and associated factors: evidence from the Nepal demographic and health survey (2001-2016). *Internat J Environ Res Pub Heal* 16: 1241.
- 7 Lamichhane R, Zhao Y, Paudel S, Adewuyi EO (2017) Factors associated with infant mortality in Nepal: a comparative analysis of Nepal demographic and health surveys (NDHS) 2006 and 2011. *BMC Publ Heal* 17: 1-18.
- 8 Acharya LB, Cleland J (2000) Maternal and child health services in rural Nepal: does access or quality matter more?. *Heal Policy Plann* 15: 223-229.
- 9 Spiegel DA, Shrestha OP, Rajbhandary T, Bijukachhe B, Sitoula P, et al. (2010) Epidemiology of surgical admissions to a children's disability hospital in Nepal. *World J Surg* 34: 954-962.
- 10 Catre D, Lopes MF, Madrigal A, Oliveiros B, Viana JS, et al. (2013) Early mortality after neonatal surgery: analysis of risk factors in an optimized health care system for the surgical newborn. *Revista Brasileira de Epidemiol* 16: 943-952.
- 11 Nagarajan N, Gupta S, Shresthra S, Varadaraj V, Devkota S, et al. (2015) Unmet surgical needs in children: a household survey in Nepal. *Pediat Surg Internat* 31: 389-395.
- 12 Weiser TG, Regenbogen SE, Thompson KD, Haynes AB, Lipsitz SR, et al. (2008) An estimation of the global volume of surgery: a modelling strategy based on available data. *The Lancet* 372: 139-144.
- 13 Walker IA, Obua AD, Mouton F, Ttendo S, Wilson IH (2010) Paediatric surgery and anaesthesia in south-western Uganda: a cross-sectional survey. *Bullet World Health Organization* 88: 897-906.
- 14 Grikscheit TC, Vacanti JP (2002) The history and current status of tissue engineering: the future of pediatric surgery. *J Pediatr Surg* 37: 277-288.
- 15 Driller C, Holschneider AM (2003) Training in pediatric surgery-a comparison of 24 countries in Europe and other countries around the world. *Europ J Pediatr Surg* 13: 73-80.