

World Pediatrics 2019: The pyrexia temperature never damage the cells of brain or harm the body - K. M. Yacob - Marma Heath Centre

K. M. Yacob

Marma Heath Centre, India

Introduction: As you aware, if temperature increases (Absence of fever) after All medicines for fever depend on the conviction that fits is the consequence of 41 degree Celsius temperature and it harms cells of mind and body. Simultaneously there is no proof based tests or solid diagnosing techniques to the conviction that fits and mind harm is the aftereffect of pyrexia. In pyrexia or nonattendance of fever a swooned persistent fell on the floor with oblivious state and annihilate cells of cerebrum, and fundamental fixings to become cognizant are same. At the point when sickness floods basic blood flow and energy level additionally decreases. At the point when the infection gets serious, capacity to siphon the blood to the cerebrum diminishes. Because of this synapses are harmed, so the patient may be incapacitated or may even pass on. At the point when the ailment gets serious, capacity to siphon the blood to the cerebrum diminishes. Because of this synapses are harmed. So the patient may be deadened or may even pass on. In pyrexia or nonattendance of fever, when blood stream to the mind diminishes and fits are framed. There is no other route than this to expand blood dissemination to the mind. It is a reasonable and tactful activity of mind to secure the life or organ. At that point the patient can encounter himself the power of blood flow. The patient can encounter when he stand his blood flow diminishes and when lie on the bed the blood dissemination diminishes. Other than that he can likewise encounter expanded blood dissemination when lie on the bed raise the foot higher than head.

In pyrexia or nonappearance of fever, when blood stream to the cerebrum diminishes and fits are shaped. There is no other route than this to build blood course to the cerebrum. It is a reasonable and circumspect activity of mind to secure the life or organ. The patient become cognizant before an opportunity to get diminishing the temperature of fever. At the point when the swooned tolerant lie on the floor, the vertical tallness among heart and cerebrum is diminished, blood flow expanded to mind. Self-checking techniques. At the point when the swooned quiet lie on the floor, the patient can stand straight and lie on bed then again. The patient become cognizant before an opportunity to get diminishing the temperature of fever. At the point when the swooned tolerant lie on the floor, the vertical tallness among heart and cerebrum is diminished, blood flow expanded to mind. All drugs for fever rely upon the conviction that fits is the result of 41 degree Celsius temperature and it hurts cells of psyche and body. At the same time there is no evidence based tests or strong diagnosing procedures to the conviction that fits and psyche hurt is the delayed consequence of pyrexia. Significant trimmings to pulverize neural

connections and fits can't be seen in fever. In pyrexia or nonattendance of fever a passed out patient fell on the floor with absent state and wreck cells of cerebrum, and fundamental trimmings to become discerning are same.

Exactly when contamination manufactures central blood scattering and energy level similarly reduces. The vertical stature among heart and psyche is more than one feet. Right when the ailment quits fooling around, ability to siphon the blood to the cerebrum reduces. In view of these neural connections are hurt. So the patient might be stifled or may even kick the pail.

In pyrexia or nonattendance of fever, when circulation system to the cerebrum reduces and fits are formed. There is no other course than this to assemble blood spread to the brain. It is a sensible and reasonable action of cerebrum to make sure about the life or organ. The patient become conscious before an occasion to get reducing the temperature of fever. Right when the fainted lenient lie on the floor, the vertical height among heart and cerebrum is reduced, blood stream extended to mind. Right when the fainted lenient lie on the floor, the patient can stand straight and lie on bed alternatively. Then the patient can experience himself the intensity of blood circulation. The patient can experience when he stand his blood scattering reduces and when lie on the bed the blood course decreases. Besides that he can in like manner experience extended blood stream when lie on the bed raise the foot higher than head.